

A JAVALLAT Immunerősítő gyógynövénycsomag hatását alátámasztó tudományos bizonyitékok

Barrett B, Brown R, Rakel D, Rabago D, et al. Placebo effects and the common cold: a randomized controlled trial. Ann.Fam.Med 2011;9:312-22.

Barrett B, Brown R, Rakel D, et al. Echinacea for treating the common cold: a randomized trial. Ann Intern Med 2010;153:769-77.

Barrett B. Medicinal properties of Echinacea: a critical review. Phytomedicine 2003;10:66-86.

Barrett BP, Brown RL, Locken K, et al. Treatment of the common cold with unrefined echinacea. A randomized, double-blind, placebo-controlled trial. Ann Intern Med 2002;137:939-46.

Buechi, S., Vogelin, R., von Eiff, M. M., Ramos, M., and Melzer, J. Open trial to assess aspects of safety and efficacy of a combined herbal cough syrup with ivy and thyme. *Forsch.Komplementarmed.Klass.Naturheilkd.* 2005;12(6):328-332.

Chavez ML, Chavez PI. Echinacea. Hosp Pharm 1998;33:180-8.

Di Pierro F, Rapacioli G, Ferrara T, Togni S. Use of a standardized extract from *Echinacea angustifolia* (Polinacea) for the prevention of respiratory tract infections. Altern Med Rev 2012;17:36-41.

Droebner K, Ehrhardt C, Poetter A, et al. CYSTUS052, a polyphenol-rich plant extract, exerts anti-influenza virus activity in mice. Antiviral Res 2007;76(1):1-10.

Ehrhardt C, Hrincius ER, Korte V, et al. A polyphenol rich plant extract, CYSTUS052, exerts anti influenza virus activity in cell culture without toxic side effects or the tendency to induce viral resistance. Antiviral Res 2007;76(1):38-47

El Hamsas El Youbi A, El Mansouri L, Boukhira S, et al. In Vivo Anti-Inflammatory and Analgesic Effects of Aqueous Extract of *Cistus ladanifer* L. from Morocco. Am J Ther 2016;23(6):e1554-9.

Ernst E, Marz R, Sieder C. A controlled multi-centre study of herbal versus synthetic secretolytic drugs for acute bronchitis. Phytomedicine 1997;4:287-93.

Ferrazzano GF, Cantile T, Roberto L, et al. Determination of the in vitro and in vivo antimicrobial activity on salivary Streptococci and Lactobacilli and chemical characterisation of the phenolic content of a *Plantago lanceolata* infusion. Biomed Res Int. 2015;2015:286817. doi: 10.1155/2015/286817.

Fiocchi A, Borella E, Riva E, et al. A double-blind clinical trial for the evaluation of the therapeutical effectiveness of a calf thymus derivative (Thymomodulin) in children with recurrent respiratory infections. Thymus 1986;8:331-9.

Galli L, de Martino M, Azzari C, et al. [Preventive effect of thymomodulin in recurrent respiratory infections in children]. Pediatr Med Chir 1990;12:229-32.

Giles JT, Palat CT III, Chien SH, et al. Evaluation of Echinacea for treatment of the common cold. Pharmacother 2000;20:690-7.

Giordani R, Regli P, Kaloustian J, et al. Antifungal effect of various essential oils against *Candida albicans*. Potentiation of antifungal action of amphotericin B by essential oil from *Thymus vulgaris*. Phytother Res 2004;18:990-5. .

Goel V, Lovlin R, Barton R, et al. Efficacy of a standardized echinacea preparation (Echinilin) for the treatment of the common cold: a randomized, double-blind, placebo-controlled trial. J Clin Pharm Ther 2004;29:75-83

Goel V, Lovlin R, Chang C, et al. A proprietary extract from the echinacea plant (*Echinacea purpurea*) enhances systemic immune response during a common cold. Phytother.Res 2005;19:689-94

Grimm W, Muller HH. A randomized controlled trial of the effect of fluid extract of *Echinacea purpurea* on the incidence and severity of colds and respiratory infections. Am J Med 1999;106:138-43.

Gruenwald, J., Graubaum, H. J., and Busch, R. Efficacy and tolerability of a fixed combination of thyme and primrose root in patients with acute bronchitis. A double-blind, randomized, placebo-controlled clinical trial. *Arzneimittelforschung* 2005;55(11):669-676.

Gruenwald, J., Graubaum, H. J., and Busch, R. Evaluation of the non-inferiority of a fixed combination of thyme fluid- and primrose root extract in comparison to a fixed combination of thyme fluid extract and primrose root tincture in patients with acute bronchitis. A single-blind, randomized, bi-centric clinical trial. *Arzneimittelforschung* 2006;56(8):574-581

Gunning K. Echinacea in the treatment and prevention of upper respiratory tract infections. *West J Med* 1999;171:198-200.

Helmholtz Zentrum München - German Research Center for Environmental Health. "Effects on HIV and Ebola: Cell culture experiments reveal potent antiviral activity of Cistus incanus. *ScienceDaily*, 2 February 2016.

Hersch-Martinez P, Leanos-Miranda BE, Solorzano-Santos F. Antibacterial effects of commercial essential oils over locally prevalent pathogenic strains in Mexico. *Fitoterapia* 2005;76:453-7.

"Hoheisel O, Sandberg M, Bertram S, et al. Echinagard treatment shortens the course of the common cold: a double blind, placebo-controlled clinical trial. *Eur J Clin Res* 1997;9:261-268.
"

Hoheisel O, Sandberg M, Bertram S, et al. Echinagard treatment shortens the course of the common cold: a double blind, placebo-controlled clinical trial. *Eur J Clin Res* 1997;9:261-268.

Jawad M, Schoop R, Suter A, et al. Safety and efficacy profile of Echinacea purpurea to prevent common cold episodes: A randomized, double-blind, placebo-controlled trial. *Evid Based Complement Alternat Med* 2012;2012:841315. Epub 2012 Sep 16.

Kalus U, Grigorov A, Kadecki O, et al. Cistus incanus (CYSTUS052) for treating patients with infection of the upper respiratory tract. A prospective, randomised, placebo-controlled clinical study. *Antiviral Res* 2009;84(3):267-71.

Karsch-Völk M, Barrett B, Kiefer D, Bauer R, Ardjomand-Woelkart K, Linde K. Echinacea for preventing and treating the common cold. *Cochrane Database Syst Rev*.2014;(2):CD000530. doi: 10.1002/14651858.CD000530.pub3.

Karsch-Völk M1, Barrett B2, Linde K1. Echinacea for preventing and treating the common cold. *JAMA*. 2015;313(6):618-9. doi: 10.1001/jama.2014.17145.

Kemmerich, B., Eberhardt, R., and Stammer, H. Efficacy and tolerability of a fluid extract combination of thyme herb and ivy leaves and matched placebo in adults suffering from acute bronchitis with productive cough. A prospective, double-blind, placebo-controlled clinical trial. *Arzneimittelforschung*. 2006;56(9):652-660.

Knols G, Stal PC, and Van Ree JW. Productive coughing complaints: Sirupus Thymi or Bromhexine? A double-blind randomized study. *Huisarts en Wetenschap* 1994;37:392-394.

Kohlert C, Schindler G, Marz RW, et al. Systemic availability and pharmacokinetics of thymol in humans. *J Clin Pharmacol* 2002;42:731-7..

Linde K, Barrett B, Wolkart K, et al. Echinacea for preventing and treating the common cold. *Cochrane Database Syst Rev* 2006;(1):CD000530.

Lindenmuth GF, Lindenmuth EB. The efficacy of echinacea compound herbal tea preparation on the severity and duration of upper respiratory and flu symptoms: a randomized, double-blind, placebo-controlled study. *J Altern Complement Med* 2000;6:327-34

Longo F, Lepore L, Agosti E, Panizon F. [Evaluation of the effectiveness of thymomodulin in children with recurrent respiratory infections]. *Pediatr Med Chir* 1988;10:603-7

Luettig B, Steinmuller C, Gifford GE, et al. Macrophage activation by the polysaccharide arabinogalactan isolated from plant cell cultures of Echinacea purpurea. *J Natl Cancer Inst* 1989;81:669-75.

Maiorano V, Chianese R, Fumarulo R, et al. Thymomodulin increases the depressed production of superoxide anion by alveolar macrophages in patients with chronic bronchitis. *Int J Tissue React* 1989;11:21-5.

Marzian, O. [Treatment of acute bronchitis in children and adolescents. Non-interventional postmarketing surveillance study confirms the benefit and safety of a syrup made of extracts from thyme and ivy leaves]. MMW.Fortschr.Med 6-28-2007;149(27-28 Suppl):69-74.

Melchart D, Walther E, Linde K, et al. Echinacea root extracts for the prevention of upper respiratory tract infections: a double-blind, placebo-controlled randomized trial. Arch Fam Med 1998;7:541-5.

O'Neil J, Hughes S, Lourie A, Zweifler J. Effects of echinacea on the frequency of upper respiratory tract symptoms: a randomized, double-blind, placebo-controlled trial. Ann Allergy Asthma Immunol 2008;100:384-8

Pina-Vaz C, Goncalves Rodrigues A, Pinto E, et al. Antifungal activity of Thymus oils and their major compounds. J Eur Acad Dermatol Venereol 2004;18:73-8.

Proestos C, Chorianopoulos N, Nychas GJ, Komaitis M. RP-HPLC analysis of the phenolic compounds of plant extracts. investigation of their antioxidant capacity and antimicrobial activity. J Agric Food Chem 2005;53:1190-5..

Ramsewak RS, Nair MG, Stommel M, Selanders L. In vitro antagonistic activity of monoterpenes and their mixtures against 'toe nail fungus' pathogens. Phytother Res 2003;17:376-9.

Raus K, Pleschka S, Klein P, Schoop R, Fisher P. Effect of an echinacea-based hot drink versus oseltamivir in Influenza treatment: a randomized, double-blind, double-dummy, multicenter, noninferiority clinical trial. . Curr Ther Res Clin Exp. 2015;20;77:66-72. doi: 10.1016/j.curtheres.2015.04.001.

Rebensburg S, Helfer M, Schneider M, et al. Potent in vitro antiviral activity of Cistus incanus extract against HIV and Filoviruses targets viral envelope proteins. Scientific Reports, 2016; 6: 20394. DOI: 10.1038/srep20394

Schoop R, Klein P, Suter A, Johnston SL. Echinacea in the prevention of induced rhinovirus colds: a meta-analysis. Clin Ther 2006;28:174-83.

Schulten B, Bulitta M, Ballering-Bruhl B, et al. Efficacy of Echinacea purpurea in patients with a common cold. A placebo-controlled, randomised, double-blind clinical trial. Arzneimittelforschung 2001;51:563-8..

Shah SA, Sander S, White CM, et al. Evaluation of echinacea for the prevention and treatment of the common cold: a meta-analysis. Lancet Infect Dis 2007;7:473-80.

Sperber SJ, Shah LP, Gilbert RD, et al. Echinacea purpurea for prevention of experimental rhinovirus colds. Clin Infect Dis 2004;38:1367-71

Stimpel M, Proksch A, Wagner H, et al. Macrophage activation and induction of macrophage cytotoxicity by purified polysaccharide fractions from the plant Echinacea purpurea. Infect Immun 1984;46:845-9.

Taylor JA, Weber W, Standish L, et al. Efficacy and safety of echinacea in treating upper respiratory tract infections in children: a randomized controlled trial. JAMA 2003;290:2824-30.

Thompson, K. D. Antiviral activity of Viracea against acyclovir susceptible and acyclovir resistant strains of herpes simplex virus. Antiviral Res 1998;39(1):55-61

Turner RB, Bauer R, Woelkart K, et al. An evaluation of Echinacea angustifolia in experimental rhinovirus infections. N Engl J Med 2005;353:341-8

Turner RB, Riker DK, Gangemi JD. Ineffectiveness of echinacea for prevention of experimental rhinovirus colds. Antimicrob Agents Chemother 2000;44:1708-9.

Vettori G, Lazzaro A, Mazzanti P, Cazzola P. [Prevention of recurrent respiratory infections in adults]. Minerva Med 1987;78:1281-9.

Vigo E, Cepeda A, Gualillo O, Perez-Fernandez R. In-vitro anti-inflammatory effect of Eucalyptus globulus and Thymus vulgaris: nitric oxide inhibition in J774A.1 murine macrophages. J Pharm Pharmacol 2004;56:257-63

Yale SH, Liu K. Echinacea purpurea therapy for the treatment of the common cold: a randomized, double-blind, placebo-controlled clinical trial. Arch Intern Med 2004;164:1237-41.