

## **Az JAVALLAT Inzulinrezisztencia csomag gyógynövényeinek hatását alátámasztó tudományos kutatások**

Atanasov, A. T. and Spasov, V. Inhibiting and disaggregating effect of gel-filtered *Galega officinalis* L. herbal extract on platelet aggregation. *J.Ethnopharmacol.* 2000;69(3):235-240.

Atanasov, A. T. and Spasov, V. Inhibiting effect of desalted extract from *Galega officinalis* L. on platelet aggregation. *Folia Med.(Plovdiv.)* 1999;41(1):46-50.

Atanasov, A. T. and Tchorbanov, B. Anti-platelet fraction from *Galega officinalis* L. inhibits platelet aggregation. *J.Med.Food* 2002;5(4):229-234.

Atanasov, A. T. and Tchorbanov, B. J. Antiplatelet aggregation activity of a fraction isolated from *Galega officinalis* L. *J Herbs Spices Medicinal Plant* 2002;10(2):63-71.

Atanasov, A. T., Chorbanov, B. P., and Dimitrov, B. D. Anti-aggregation activity of crude water extract of *Galega officinalis* L. fractionated on Sephadex G-25 and Sepharose 4B. *Folia Med. (Plovdiv.)* 2002;44(1-2):45-49.

Broadhurst CL, Polansky MM, Anderson RA. Insulin-like biological activity of culinary and medicinal plant aqueous extracts in vitro. *J Agric Food Chem* 2000;48:849-52..

Chrubasik C, Maier T, Dawid C, et al. An observational study and quantification of the actives in a supplement with *Sambucus nigra* and *Asparagus officinalis* used for weight reduction. *Phytother Res* 2008;22:913-8

Edgcumbe, D. P. and McAuley, D. Hypoglycaemia related to ingestion of a herbal remedy. *Eur.J.Emerg.Med.* 2008;15(4):236-237.

Eidi, M., Eidi, A., and Zamanizadeh, H. Effect of *Salvia officinalis* L. leaves on serum glucose and insulin in healthy and streptozotocin-induced diabetic rats. *J Ethnopharmacol.* 9-14-2005;100(3):310-313.

Ghalavand A, Motamedi P, Deleramnasab M, Khodadoust M. The effect of interval training and nettle supplement on glycemic control and blood pressure in men with Type 2 Diabetes. *Int J Basic Sci Med.* 2017;2(1):33-40.

Gray AM, Abdel-Wahab YH, Flatt PR. The traditional plant treatment, *Sambucus nigra* (elder), exhibits insulin-like and insulin-releasing actions in vitro. *J Nutr* 2000;130:15-20.  
Khalili N, Fereydoonzadeh R, Mohtashami R, Mehrzadi S, Heydari M, Huseini F. Silymarin, Olibanum, and Nettle, a mixed herbal formulation in the treatment of Type II Diabetes: a randomized, double-blind, placebo-controlled, clinical trial. *J Evid Based Complementary Altern Med.* 2017 Oct;22(4):603-608

Kianbakht S, Dabaghian FH. Improved glycemic control and lipid profile in hyperlipidemic type 2 diabetic patients consuming *Salvia officinalis* L. leaf extract: a randomized placebo. Controlled clinical trial. *Complement Ther Med.* 2013;21(5):441-6.

Kianbakht S, Khalighi-Sigaroodi F, Dabaghian FH. Improved glycemic control in patients with advanced type 2 diabetes mellitus taking *Urtica dioica* leaf extract: a randomized double-blind placebo-controlled clinical trial. *Clin. Lab.* 2013;59:1071-1076.

Kwon, Y. I., Vattem, D. A., and Shetty, K. Evaluation of clonal herbs of Lamiaceae species for management of diabetes and hypertension. *Asia Pac.J Clin Nutr* 2006;15(1):107-118  
Lemus, I., Garcia, R., Delvillar, E., and Knop, G. Hypoglycaemic activity of four plants used in Chilean popular medicine. *Phytother Res* 1999;13(2):91-94

Mooney, M. H., Fogarty, S., Stevenson, C., Gallagher, A. M., Palit, P., Hawley, S. A., Hardie, D. G., Coxon, G. D., Waigh, R. D., Tate, R. J., Harvey, A. L., and Furman, B. L. Mechanisms underlying the metabolic actions of galegine that contribute to weight loss in mice. *Br.J Pharmacol.* 2008;153(8):1669-1677.

Neef, H., Augustijns, P., Declercq, P., Declerck, P. J., and Laekeman, G. Inhibitory effects of *Galega officinalis* on glucose transport across monolayers of human intestinal epithelial cells. *Pharmaceutical and Pharmacological Letters* 1996;6:86-89.

Palit, P., Furman, B. L., and Gray, A. I. Novel weight-reducing activity of *Galega officinalis* in mice. *J Pharm Pharmacol.* 1999;51(11):1313-1319.

Perry, N. S., Houghton, P. J., Sampson, J., Theobald, A. E., Hart, S., Lis-Balchin, M., Hoult, J. R., Evans, P., Jenner, P., Milligan, S., and Perry, E. K. In-vitro activity of *S. lavandulaefolia* (Spanish sage) relevant to treatment of Alzheimer's disease. *J Pharm Pharmacol* 2001;53(10):1347-1356.

Petricic, J. and Kalodera, Z. Galegin in the goat's rue herb: toxicity, antidiabetic activity and content determination. *Acta Pharm Jugosl* 1982;32:219-223.

Rau, O., Wurglits, M., Paulke, A., Zitzkowski, J., Meindl, N., Bock, A., Dingermann, T., Abdel-Tawab, M., and Schubert-Zsilavecz, M. Carnosic Acid and Carnosol, Phenolic Diterpene Compounds of the Labiate Herbs Rosemary and Sage, are Activators of the Human Peroxisome Proliferator-Activated Receptor Gamma. *Planta Med* 2006;72(10):881-887.

Viapiana A, Wesolowski M. The phenolic contents and antioxidant activities of infusions of *Sambucus nigra* L. *Plant Foods Hum Nutr.* 2017;72(1):82-7.

Ziae R, Foshati S, Hadi A, et al. The effect of nettle (*Urtica dioica*) supplementation on the glycemic control of patients with type 2 diabetes mellitus: a systematic review and meta-analysis. *Phytother Res* 2020;34(2):282-94.