

## A JAVALLAT Diabétesz gyógynövénycsomag hatását alátámasztó tudományos bizonyítékok

Alarcon-Aguilar FJ, Roman-Ramos R, Flores-Saenz JL, Aguirre-Garcia F. Investigation on the hypoglycaemic effects of extracts of four Mexican medicinal plants in normal and alloxan-diabetic mice. *Phytother Res.* 2002;16(4):383-6.

Amini L, Mojab F, Jahanfar S, Sepidarkish M, Raoofi Z, Maleki-Hajiaghah A. Efficacy of *Salvia officinalis* extract on the prevention of insulin resistance in euglycemic patients with polycystic ovary syndrome: A double-blinded placebo-controlled clinical trial. *Complement Ther Med.* 2020;48:102245

Bakhshayeshi S, Madani SP, Hemmatabadi M, et al. Effects of Semelil (ANGIPARS) on diabetic peripheral neuropathy: a randomized, double-blind placebo-controlled clinical trial. *Daru: Journal of Faculty of Pharmacy, Tehran University of Medical Sciences.* 2011;19(1):65.

Boivin M, Flourie B, Rizza RA, Go VL, DiMagno EP. Gastrointestinal and metabolic effects of amylase inhibition in diabetics. *Gastroenterology.* 1988;94(2):387-94.

Cheang KI, Nguyen TT, Karjane NW, Salley KE. Raspberry Leaf and Hypoglycemia in Gestational Diabetes Mellitus. *Obstet Gynecol.* 2016;128(6):1421-4

Eidi, M., Eidi, A., and Zamanizadeh, H. Effect of *Salvia officinalis* L. leaves on serum glucose and insulin in healthy and streptozotocin-induced diabetic rats. *J Ethnopharmacol.* 9-14-2005;100(3):310-313.

Hamidpour M, Hamidpour R, Hamidpour S, Shahlari M. Chemistry, Pharmacology, and Medicinal Property of Sage (*Salvia*) to Prevent and Cure Illnesses such as Obesity, Diabetes, Depression, Dementia, Lupus, Autism, Heart Disease, and Cancer. *J Tradit Complement Med.* 2014;4(2):82-8.

Ionescu-Tirgoviste, C., Popa, E., Mirodon, Z., Simionescu, M., and Mincu, I. [The effect of a plant mixture on the metabolic equilibrium in patients with type-2 diabetes mellitus]. *Rev.Med Interna Neurol.Psihiatr.Neurochir.Dermatovenerol.Med Interna* 1989;41(2):185-192.

Kianbakht S, Dabaghian FH. Improved glycemic control and lipid profile in hyperlipidemic type 2 diabetic patients consuming *Salvia officinalis* L. leaf extract: a randomized placebo. Controlled clinical trial. *Complement Ther Med.* 2013;21(5):441-6.

Kwon, Y. I., Vattem, D. A., and Shetty, K. Evaluation of clonal herbs of Lamiaceae species for management of diabetes and hypertension. *Asia Pac.J Clin Nutr* 2006;15(1):107-118.

Lima, C. F., Azevedo, M. F., Araujo, R., Fernandes-Ferreira, M., and Pereira-Wilson, C. Metformin-like effect of *Salvia officinalis* (common sage): is it useful in diabetes prevention? *Br.J Nutr* 2006;96(2):326-333.

Olmedilla-Alonso B, Pedrosa MM, Cuadrado C, et al. Composition of two Spanish common dry beans (*Phaseolus vulgaris*), 'Almonga' and 'Curruquilla', and their postprandial effect in type 2 diabetics. *J Sci Food Agric* 2013;93(5):1076-82.

Rondanelli M, Giacosa A, Orsini F, Opizzi A, Villani S. Appetite control and glycaemia reduction in overweight subjects treated with a combination of two highly standardized extracts from *Phaseolus vulgaris* and *Cynara scolymus*. *Phytother Res* 2011 Sep;25(9):1275-82.

Spadafranca A, Rinelli S, Riva A, et al. *Phaseolus vulgaris* extract affects glycometabolic and appetite control in healthy human subjects. *Br J Nutr* 2013;109(10):1789-95.

Tjokroprawiro A, Pikir BS, Budhiarta AA, et al. Metabolic effects of onion and green beans on diabetic patients. *Tohoku J Exp Med* 1983;141 Suppl:671-6

Venkateswaran, S., Pari, L., and Saravanan, G. Effect of *Phaseolus vulgaris* on circulatory antioxidants and lipids in rats with streptozotocin-induced diabetes. *J Med Food* 2002;5(2):97-103.

Winham DM, Hutchins AM, Melde CL. Pinto bean, navy bean, and black-eyed pea consumption do not significantly lower the glycemic response to a high glycemic index treatment in normoglycemic adults. *Nutrition Res* 2007;27(9):535-541.